

Multiwavelength observations of ULXs

Mar Mezcuá

University of Montréal



Ultraluminous X-ray sources (ULXs)

- Extragalactic, off-nuclear, point-like X-ray sources with $L_X > L_{\text{Edd}}$ for a stellar-mass black hole:

$$L_{\text{Edd}} = 1.3 \times 10^{38} M_{\text{BH}}/M_{\odot} \text{ erg/s}$$

Ultraluminous X-ray sources (ULXs)

- Extragalactic, off-nuclear, point-like X-ray sources with $L_X > L_{\text{Edd}}$ for a stellar-mass black hole:

$$L_{\text{Edd}} = 1.3 \times 10^{38} M_{\text{BH}}/M_{\odot} \text{ erg/s}$$

$$L_X \geq 10^{39} \text{ erg/s for a } 10 M_{\odot} \text{ BH}$$




Ultraluminous X-ray sources (ULXs)

- Extragalactic, off-nuclear, point-like X-ray sources with $L_X > L_{\text{Edd}}$ for a stellar-mass black hole:

$$L_{\text{Edd}} = 1.3 \times 10^{38} M_{\text{BH}}/M_{\odot} \text{ erg/s}$$

$$L_X \geq 10^{39} \text{ erg/s for a } 10 M_{\odot} \text{ BH}$$

- Most ULXs: $L_X \sim 10^{39} - 10^{40} \text{ erg/s}$
Extreme ULXs: $L_X > 2 \times 10^{40} \text{ erg/s}$
Hyperluminous X-ray sources (HLXs): $L_X \geq 10^{41} \text{ erg/s}$



NASA Chandra, HST, Spitzer

The nature of ULXs

Is the X-ray emission isotropic?

No, it's beamed!

Yes, it is!

Is it a black hole?

Relativistic
beaming

Mild beaming

Yes, it's a BH!

No!

super-Eddington
accretion

sub-Eddington
accretion

Microquasar

Stellar-mass
BH

Intermediate-
mass BH

Neutron
star

The nature of ULXs

Is the X-ray source

No, it's beamed!

Relativistic
beaming

Mild beaming

Microquasar



Is it a black hole?

Is a BH?

Sub-Eddington
accretion

No!

Stellar-mass
BH

Intermediate-
mass BH

Neutron
star

CHECKLIST

- ☐ Spectral analysis
- ☐ Variability
- ☐ Timing analysis
- ☐ Distance
- ☐ Mass measurement
- ☐ Environment
- ☐ Jets/outflows
- ☐ Monitoring

CHECKLIST

☒ Spectral analysis

☒ Variability

☒ Timing analysis

☐ Distance

☐ Mass measurement

☐ Environment

☐ Jets/outflows

☐ Monitoring

X-rays

CHECKLIST

☒ Spectral analysis

☒ Variability

☒ Timing analysis

☒ Distance

☒ Mass measurement

☒ Environment

☒ Jets/outflows

☒ Monitoring


X-rays

Optical, IR
radio

✓ Spectral analysis

Most ULXs spectra:

- soft excess at low energies
- power-law cutoff above ~ 3 keV



Bachetti et al. (2013)

Also e.g., Stobbart et al. (2006),


Walton et al. (2011, 2013, 2014, 2015),

Rana et al. (2015)

✓ Spectral analysis

Most ULXs spectra:

- soft excess at low energies
- power-law cutoff above ~ 3 keV



Characteristic of
'ultraluminous' state
Interpreted as
super-Eddington
accretion

Gladstone et al. (2009)
Sutton et al. (2013)


Bachetti et al. (2013)
Also e.g., Stobbart et al. (2006),
Walton et al. (2011, 2013, 2014, 2015),
Rana et al. (2015)

✓ Variability

- Short-term variability observed in many ULXs
- Explained by super-Eddington wind

Middleton et al. (2011)

More by Matt Middleton!



Also e.g., King (2004, 2009)
Poutanen et al. (2007)
King & Pounds (2003)
Middleton et al. (2014, 2015)
Sutton et al. (2013)

✓ Variability


- Short-term variability observed in many ULXs
- Explained by super-Eddington wind

Middleton et al. (2011)

More by Matt Middleton!

- Observational evidence of outflow wind at $0.2c$!!


Talk by Ciro Pinto!



Also e.g., King (2004, 2009)
Poutanen et al. (2007)
King & Pounds (2003)
Middleton et al. (2014, 2015)
Sutton et al. (2013)

✓ Timing analysis



- Twin-peak high-frequency QPOs found in some ULXs



M82 X-1

Pasham et al. (2014, 2015)


NGC 1313 X-1



✓ Timing analysis

- Twin-peak high-frequency QPOs found in some ULXs
- Frequency $\sim 1/M_{\text{BH}}$ in stellar-mass BHs



Extrapolating... $\Rightarrow$ **IMBH!**



M82 X-1


Pasham et al. (2014, 2015)

NGC 1313 X-1



✓ Timing analysis

- Low-frequency QPOs found in some ULXs
- Frequency scaling from stellar-mass BHs $\Rightarrow$ **IMBH!**




M82 X-2

Feng et al. (2010)

✓ Timing analysis


- Low-frequency QPOs found in some ULXs
- Frequency scaling from stellar-mass BHs $\Rightarrow$ **IMBH!**



M82 X-2

Feng et al. (2010)

But...pulsations reveal neutron star!



Bachetti et al. (2014)

CHECKLIST

- ☒ Spectral analysis
- ☒ Variability
- ☒ Timing analysis
- ☒ Distance
- ☒ Mass measurement
- ☒ Environment
- ☒ Jets/outflows
- ☒ Monitoring

X-rays

Optical, IR
radio

CHECKLIST

- ☒ Spectral analysis
- ☒ Variability
- ☒ Timing analysis
- ☒ Distance
- ☒ Mass measurement
- ☒ Environment
- ☒ Jets/outflows
- ☒ Monitoring

X-rays

Optical, IR
radio

Optical counterparts



- Tens of ULXs with optical counterpart

Thanks to Chandra+HST astrometry

- Emission tends to be blue

$m_V \sim 20-26$

Gladstone et al. (2013)



Optical counterparts

- Tens of ULXs with optical counterpart

Thanks to Chandra+HST astrometry

- Emission tends to be blue

$m_V \sim 20-26$

BUT optical emission from:



?

?

?

?

Gladstone et al. (2013)



Optical counterparts

- Tens of ULXs with optical counterpart

Thanks to Chandra+HST astrometry



- Emission tends to be blue

$m_V \sim 20-26$

BUT optical emission from:

- ? * companion star? ?
- * intrinsic accretion disk?
- ? * X-ray irradiated outer disk?
- * all the above combined? ?

Gladstone et al. (2013)




HST data of 13 ULXs:

- SED: no single spectral type match
- SED: power-law fit $\alpha \sim 1-2$ ($F_\nu \sim \nu^\alpha$)
- optical variability
- $\log(f_x/f_v) \sim 3$ consistent with LMXBs



optical emission dominated by
X-ray reprocessing on the outer disk




Tao et al. (2011)

HST data of 13 ULXs:

- SED: no single spectral type match
- SED: power-law fit $\alpha \sim 1-2$ ($F_\nu \sim \nu^\alpha$)
- optical variability
- $\log(f_x/f_v) \sim 3$ consistent with LMXBs



optical emission dominated by
X-ray reprocessing on the outer disk



Tao et al. (2011)

Mmmm...
What could we do...?



Apply irradiation models!

X-ray binary

Roche lobe
overflow

Parameters: mass transfer rate,
BH mass, inclination, donor
radius, donor mass...

stellar/binary evolutionary tracks



e.g., Copperwheat et al. (2005, 2007), Patruno & Zampieri (2008, 2010),
Madhusudhan et al. (2008), Grisé et al. (2011), Gladstone et al. (2013)


Apply irradiation models!

X-ray binary

Roche lobe
overflow

Parameters: mass transfer rate,
BH mass, inclination, donor
radius, donor mass...

stellar/binary evolutionary tracks



Reproduce optical colors
Predict optical luminosity
of emission from star &
accretion disk

e.g., Copperwheat et al. (2005, 2007), Patruno & Zampieri (2008, 2010),
Madhusudhan et al. (2008), Grisé et al. (2011), Gladstone et al. (2013)


Apply irradiation models!

X-ray binary

Roche lobe
overflow

Parameters: mass transfer rate,
BH mass, inclination, donor
radius, donor mass...

stellar/binary evolutionary tracks



Most donor
stars consistent
with B giants

e.g., Copperwheat et al. (2005, 2007), Patruno & Zampieri (2008, 2010),
Madhusudhan et al. (2008), Grisé et al. (2011), Gladstone et al. (2013)

☑ Distance

Optical counterpart or background AGN?

Spectroscopy needed!

10 ULXs: 2 background AGN

Sutton et al. (2015)

17 ULXs: 15 background AGN
2 foreground stars


Wong et al. (2008)

23 ULXs: 20 background AGN
3 foreground stars

Gutiérrez & López-Corredoira (2005, 2007)

Gutiérrez (2006, 2013)

Sutton et al. (2015)



HLX in IC 4320

$m_V = 22.67$

☑ Distance

Optical counterpart or background AGN?

Spectroscopy needed!

10 ULXs: 2 background AGN

Sutton et al. (2015)

17 ULXs: 15 background AGN
2 foreground stars

Wong et al. (2008)

23 ULXs: 20 background AGN
3 foreground stars

Gutiérrez & López-Corredoira (2005, 2007)

Gutiérrez (2006, 2013)

Sutton et al. (2015)



HLX in IC 4320

$m_V = 22.67$

☑ Distance

Optical counterpart or background AGN?

Spectroscopy needed!

10 ULXs: 2 background AGN

Sutton et al. (2015)

17 ULXs: 15 background AGN
2 foreground stars

Wong et al. (2008)

23 ULXs: 20 background AGN
3 foreground stars

Gutiérrez & López-Corredoira (2005, 2007)

Gutiérrez (2006, 2013)



HLX in IC 4320

$m_V = 22.67$

ULX location confirmed for many other ULXs

e.g., Pakull & Mirioni (2002), Lehmann et al. (2005), Grisé et al. (2006), Pakull, Grisé & Motch (2006), Kaaret & Corbel (2009), Wiersema et al. (2010), Motch et al. (2011), Roberts et al. (2011), Soria et al. (2013), Gutiérrez & Moon (2014)

Meanwhile, in the infrared...

Spectroscopy:

- 4 ULXs: 3 background AGN
- $\log(f_x/f_v) = -1 - 1 \Rightarrow$ Likely AGN

Heida et al. (2013)

Meanwhile, in the infrared...

Spectroscopy:

- 4 ULXs: 3 background AGN
- $\log(f_x/f_v) = -1 - 1 \Rightarrow$ Likely AGN

Heida et al. (2013)

Photometry & spectroscopy:

- ULX counterparts consistent with **red supergiants**
- radial velocity confirms ULX location for a few ULXs
- but no dynamical mass measurement.... YET!

★ Twinkle, twinkle ★
★ ~~little~~ star... ★

Heida et al. (2014, 2015, 2016)

Meanwhile, in the infrared...

Spectroscopy:

- 4 ULXs: 3 background AGN
- $\log(f_x/f_v) = -1 - 1 \Rightarrow$ Likely AGN

Heida et al. (2013)

Photometry & spectroscopy:

- ULX counterparts consistent with **red supergiants**
- radial velocity confirms ULX location for a few ULXs
- but no dynamical mass measurement.... YET!

★ Twinkle, twinkle ★
★ ~~little~~ star... ★

Heida et al. (2014, 2015, 2016)

See next talks by Marianne
Heida & Kristhell López!

✓ Dynamical mass measurement

Several attempts in the optical...

e.g., Pakull et al. (2006), Kaaret & Corbel (2009), Pakull, Grisé & Motch (2006), Grisé et al. (2009), Roberts et al. (2011), Liu et al. (2012)

• M101 X-1

$L_x = 3 \times 10^{39}$ erg/s, Wolf-Rayet donor of $19 M_\odot$

✓ Dynamical mass measurement

Several attempts in the optical...

e.g., Pakull et al. (2006), Kaaret & Corbel (2009), Pakull, Grisé & Motch (2006), Grisé et al. (2009), Roberts et al. (2011), Liu et al. (2012)

• M101 X-1

$L_X = 3 \times 10^{39}$ erg/s, Wolf-Rayet donor of $19 M_\odot$

Liu et al. (2011)


Radial velocity variations:

± 60 km/s over 3 months

Orbital period: 8.2 ± 0.1 days



$5 < M_{BH} < 20 M_\odot$
super-Eddington accretion



✓ Dynamical mass measurement

Several attempts in the optical...

e.g., Pakull et al. (2006), Kaaret & Corbel (2009), Pakull, Grisé & Motch (2006), Grisé et al. (2009), Roberts et al. (2011), Liu et al. (2012)

• M101 X-1

$L_X = 3 \times 10^{39}$ erg/s, Wolf-Rayet donor of $19 M_\odot$

Liu et al. (2011)

Radial velocity variations:

± 60 km/s over 3 months


Orbital period: 8.2 ± 0.1 days



$5 < M_{BH} < 20 M_\odot$
super-Eddington accretion

Signatures of optical thick outflow!

Shen et al. (2015), Soria & Kong (2016)




• ULX P13 in NGC 7793

Donor star B9Ia

Optical/UV modulations

Orbital period ~ 64 days

 $M_{\text{BH}} < 15 M_{\odot}$
super-Eddington accretion

ULX P13 in NGC 7793


Donor star B9Ia

Optical/UV modulations

Orbital period ~ 64 days



$M_{\text{BH}} < 15 M_{\odot}$
super-Eddington accretion




Motch et al (2014)

$L_x \sim 4 \times 10^{39} \text{ erg/s}$

Soft excess


Spectral break at ~ 4.2 keV

Signatures of
super-Eddington accretion



✓ Environment

- Some ULXs near/within >10 Myr old stellar clusters
- Others near star formation ring




Kaaret (2005)



Grisé et al. (2011)

e.g., Liu et al. (2007), Feng & Kaaret (2008), Grisé et al. (2008, 2011), Yang et al. (2011), Farrell et al. (2012)



- Some ULXs surrounded by nebular emission



Feng & Soria (2011)


Shock-ionized
nebulae

X-ray photoionized nebulae



e.g., Pakull & Mirioni (2002, 2003), Pakull & Grisé (2008), Abolmasov et al. (2007), Kaaret & Corbel (2009), Kaaret et al. (2004, 2010), Moon et al. (2011), Cseh et al. (2012)

- Some ULXs surrounded by nebular emission





Feng & Soria (2011)

Shock-ionized nebulae


e.g., Pakull & Mirioni (2002, 2003), Pakull & Grisé (2008), Abolmasov et al. (2007), Kaaret & Corbel (2009), Kaaret et al. (2004, 2010), Moon et al. (2011), Cseh et al. (2012)

X-ray photoionized nebulae

$L_x \sim 10^{39} - 10^{40}$ erg/s required to produce high-ionization (e.g., HeII) line luminosities



- Some ULXs surrounded by nebular emission



Feng & Soria (2011)

Shock-ionized nebulae



e.g., Pakull & Mirioni (2002, 2003), Pakull & Grisé (2008), Abolmasov et al. (2007), Kaaret & Corbel (2009), Kaaret et al. (2004, 2010), Moon et al. (2011), Cseh et al. (2012)

X-ray photoionized nebulae


$L_x \sim 10^{39} - 10^{40}$ erg/s required to produce high-ionization (e.g., HeII) line luminosities



Isotropic X-ray emission




- Also radio nebulae




Optically-thin synchrotron emission

Size ~ 40-200 pc

More luminous than SNR



- Also radio nebulae



Optically-thin synchrotron emission


Size ~ 40-200 pc

More luminous than SNR


"ULX bubbles"

Shock-ionized nebulae

Radio nebulae



- Also radio nebulae



Optically-thin synchrotron emission

Size $\sim 40\text{--}200$ pc

More luminous than SNR

"ULX bubbles"


Shock-ionized nebulae
Radio nebulae




Inflated by jets or winds

Energy carried $\sim 10^{49} - 10^{53}$ erg/s

(SS433 $\sim 10^{46}$ erg/s)




- Radio jets inside nebula



Radio bubble inflated by jet
(although optical nebula
produced by photoionization!)


Cseh et al. (2014, 2015)

- Radio jets inside nebula




Radio bubble inflated by jet
(although optical nebula
produced by photoionization!)

Cseh et al. (2014, 2015)




Optically thin synchrotron emission
Has faded factor ~ 7 in 1.5 yrs
Has cooled faster than outer components

- Radio jets inside nebula



Radio bubble inflated by jet
(although optical nebula
produced by photoionization!)

Cseh et al. (2014, 2015)



Optically thin synchrotron emission
Has faded factor ~ 7 in 1.5 yrs
Has cooled faster than outer components



Radio bubble inflated by jet EJECTA

• Radio flares

HLX-1

$$L_x \sim 10^{42} \text{ erg/s}$$

Periodic outbursts ~months



Spectral state transitions ~day

Transient jet radio emission follows
X-ray spectral state transitions
(as in Galactic XRBs!)

$$M_{\text{BH}} \sim 10^4 - 10^5 M_{\odot} \Rightarrow \text{IMBH}$$

Nucleus of stripped dwarf galaxy?

e.g., Farrell et al. (2009), Davis et al. (2011), Lasota (2011), Servillat et al. (2011), Godet et al. (2009, 2012, 2014), Yan et al. (2015), Cseh et al. (2015)



Webb et al. (2012)

XMMU J004243.6+412519 in M31



$L_X \sim 2 \times 10^{38} \text{ erg/s} \rightarrow$ sub-Eddington regime

$L_X \sim 1.2 \times 10^{39} \text{ erg/s} \rightarrow$ ULX, super-Eddington regime

Radio flares, similar behavior as Galactic XRBs

Radio variability ~days & minutes $\rightarrow$ size $\sim 5 \text{ AU}$

Middleton et al. (2013)



- Compact radio emission

NGC 5457 X-9

$$L_x \sim 3 \times 10^{39} \text{ erg/s}$$

Short- & long-term X-ray variability

$$\Gamma \sim 0.2 - 0.4 \text{ or cool disc?}$$


Jenkins et al. (2004, 2005)

Fundamental plane of BH accretion
(if sub-Eddington):



IMBH

Mezcua et al. (2013a)





NGC 2276-3c

$L_x \sim 6 \times 10^{40} \text{ erg/s}$

Short-term X-ray variability

$\Gamma \sim 1.4$



Sutton et al. (2012)

Mezcua et al. (2013c, 2015)

NGC 2276-3c

$$L_x \sim 6 \times 10^{40} \text{ erg/s}$$

Short-term X-ray variability


$$\Gamma \sim 1.4$$

Radio jet $\sim 2 \text{ pc}$, flat α



Fundamental plane



$$M_{\text{BH}} \sim 5 \times 10^4 M_{\odot}$$



HST



Sutton et al. (2012)

Mezcua et al. (2013c, 2015)

Summary

ULXs: mixed bag of objects



- X-rays: soft excess, high-energy cut-off
- Optical: dynamical mass measurements

stellar-mass BHs
super-Eddington
accretion

Summary

ULXs: mixed bag of objects



- X-rays: soft excess, high-energy cut-off
- Optical: dynamical mass measurements
- X-rays: high-frequency QPOs
- Radio: jets, fundamental plane



stellar-mass BHs
super-Eddington
accretion



IMBHs

Summary

ULXs: mixed bag of objects



- X-rays: soft excess, high-energy cut-off
- Optical: dynamical mass measurements
- X-rays: high-frequency QPOs
- Radio: jets, fundamental plane
- X-rays: pulsations

stellar-mass BHs
super-Eddington
accretion

IMBHs

neutron star

Summary

ULXs: mixed bag of objects



- X-rays: soft excess, high-energy cut-off
- Optical: dynamical mass measurements
- X-rays: high-frequency QPOs
- Radio: jets, fundamental plane
- X-rays: pulsations
- Optical/IR spectroscopy: background AGN?
- Radio/optical: environment, nebulae
- Radio/X-ray monitoring: similarities with XRBs

stellar-mass BHs
super-Eddington
accretion

IMBHs

neutron star

THANK YOU!!



www.astro.umontreal.ca/~marmezcua/



@marmezcua_astro